

PRE SURGICAL TESTING REQUIREMENTS

HISTORY AND PHYSICAL

All Patients Within 30 days of surgery

EKG

Any patient with Diabetes, Hypertension, Cardiac, Vascular, Pulmonary, Renal, or Hepatic Disease

All Men > 45 years old

All Women > 55 years old

Within 3 months of surgery

CHEST X-RAY

Not required

LABORATORY WORK

Within 30 days of surgery

	<u>General Anesthesia</u>	<u>MAC</u>	<u>Anterior Segment Surgery under MAC only</u>
Healthy Patient	none	none	none
Diabetes Hypertension Cardiac/Pulmonary Renal	BMP	BMP	none
Liver disease	CBC, BMP PT/PTT, LFT	CBC, BMP PT/PTT	none
Coumadin therapy	INR	INR	none

For history of anemia or for surgeries where blood loss is expected to be >200cc, please include CBC

For patients on kidney dialysis, K+ should be obtained day of surgery

All diabetic patients glucose levels (i.e. finger stick) to be checked day of surgery

Urine pregnancy day of admission for all women of menstruating age

For patients with AICDs, please see NYEE's policy concerning defibrillators

Patients with more complex medical conditions may require further workup (i.e stress tests, echocardiogram, cardio/pulmonary consult, etc). Please consult anesthesia department or patient's PMD.

Cataract Surgery under MAC does not require an EKG if there are no coexisting diseases, regardless of age.

CBC = complete blood count, BMP = basic metabolic profile, LFT = liver ftinction test, K+ = potassium

PT/PTT/INR = prothrombin time/partial prothrombin time/international normalized ratio

AICD = internal cardiac defibrillator